

AD-A169 100


ON THE BOUNDED REGRET OF EMPIRICAL BAYES ESTIMATORS(U)
SOUTH CAROLINA UNIV COLUMBIA DEPT OF STATISTICS K F YU
MAY 85 TR-112 AFOSR-TR-86-0276 AFOSR-84-0156


1/1

UNCLASSIFIED

F/B 12/1

NL





W. R. H. 1.0 1.1 1.25 1.4 1.6 1.8 2.0 2.2 2.5 2.8 3.2 3.6 4.0 4.5

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

(2)

REPORT DOCUMENT

AD-A169 108

1a. REPORT SECURITY CLASSIFICATION
UNCLASSIFIED

2a. SECURITY CLASSIFICATION AUTHORITY

2b. DECLASSIFICATION/DOWNGRADING SCHEDULE

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

3. DISTRIBUTION/AVAILABILITY OF REPORT

Approved for public release; distribution unlimited.

5. MONITORING ORGANIZATION REPORT NUMBER(S)

AFOSR-TR- 86-0276

6a. NAME OF PERFORMING ORGANIZATION

University of South Carolina

6b. OFFICE SYMBOL
(If applicable)

7a. NAME OF MONITORING ORGANIZATION

Air Force Office of Scientific Research

6c. ADDRESS (City, State and ZIP Code)

Department of Statistics
Statistical Laboratory
Columbia S.C. 29208

7b. ADDRESS (City, State and ZIP Code)

Directorate of Mathematical & Information
Sciences, Bolling AFB DC 20332-64488a. NAME OF FUNDING/SPONSORING
ORGANIZATION

AFOSR

8b. OFFICE SYMBOL
(If applicable)

NM

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

AFOSR-84-0156

8c. ADDRESS (City, State and ZIP Code)

Bolling AFB DC 20332-6448

10. SOURCE OF FUNDING NOS.

PROGRAM
ELEMENT NO.
61102FPROJECT
NO.
2304TASK
NO.
A5WORK UNIT
NO.

11. TITLE (Include Security Classification)

ON THE BOUNDED REGRET OF EMPIRICAL BAYES ESTIMATORS

12. PERSONAL AUTHOR(S)

Kai F. Yu

13a. TYPE OF REPORT

Technical

13b. TIME COVERED

FROM TO

14. DATE OF REPORT (Yr., Mo., Day)

May 85

15. PAGE COUNT

14

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD GROUP SUB GR

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

ABSTRACT

Let $(\theta_1, x_1), \dots, (\theta_n, x_n)$ be independent and identically distributed random vectors with $E(x|\theta) = \theta$ and $\text{Var}(x|\theta) = a + b\theta + c\theta^2$. Let t_1 be the linear Bayes estimator of θ_1 and $\tilde{\theta}_1$ be the linear empirical Bayes estimator of θ_1 as proposed in Robbins (1983), when $E x$ and $\text{Var } x$ are unknown to the statistician. The regret of using $\tilde{\theta}_1$ instead of t_1 because of ignorance of the mean and the variance is $r_1 = E(\tilde{\theta}_1 - \theta_1)^2 - E(t_1 - \theta_1)^2$.

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS ☐

21. ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

22a. NAME OF RESPONSIBLE INDIVIDUAL

Major Brian W. Woodruff

22b. TELEPHONE NUMBER
(Include Area Code)

(202) 767- 5027

22c. OFFICE SYMBOL

NM

DTIC FILE COPY

DTIC
ELECTE
JUN 10 1986
S D

86 6 10 149

Under appropriate conditions cumulative regret $R_n = r_1 + \dots + r_n$ is shown to have a finite limit even when n tends to infinity. The limit can be explicitly computed in terms of a, b, c and the first four moments of x .

AFOSR-TR- 36 - 0276

ON THE BOUNDED REGRET OF EMPIRICAL BAYES ESTIMATORS

by

Kai F. Yu

University of South Carolina
Statistics Technical Report No. 112
62C12-1

DEPARTMENT OF STATISTICS

**The University of South Carolina
Columbia, South Carolina 29208**

Approved for public release;
distribution unlimited.

2

ON THE BOUNDED REGRET OF EMPIRICAL BAYES ESTIMATORS

by

Kai F. Yu

University of South Carolina
Statistics Technical Report No. 112
62C12-1

May, 1985

Department of Statistics
University of South Carolina
Columbia, SC 29208

DTIC
ELECTE
JUN 10 1986
S D D

AIR FORCE RESEARCH AND DEVELOPMENT COMMAND
AFRL-ML-85-001
1. Title
2. Author
3. Report Number
4. Date
5. Distribution
6. Notes

ON THE BOUNDED REGRET OF EMPIRICAL BAYES ESTIMATORS

Kai F. Yu

Department of Statistics
University of South Carolina
Columbia, SC 29208

Key Words and Phrases: Empirical Bayes; bounded regret; uniform integrability.

ABSTRACT

Let $(\theta_1, x_1), \dots, (\theta_n, x_n)$ be independent and identically distributed random vectors with $E(x|\theta) = \theta$ and $\text{Var}(x|\theta) = a + b\theta + c\theta^2$. Let t_i be the linear Bayes estimator of θ_i and $\tilde{\theta}_i$ be the linear empirical Bayes estimator of θ_i as proposed in Robbins (1983), when $E x$ and $\text{Var } x$ are unknown to the statistician. The regret of using $\tilde{\theta}_i$ instead of t_i because of ignorance of the mean and the variance is $r_i = E(\tilde{\theta}_i - \theta_i)^2 - E(t_i - \theta_i)^2$. Under appropriate conditions cumulative regret $R_n = r_1 + \dots + r_n$ is shown to have a finite limit even when n tends to infinity. The limit can be explicitly computed in terms of a, b, c and the first four moments of x .

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Dist	Avail and/or S. Avail
A-1	



1. INTRODUCTION AND SUMMARY

In the first Jerzy Neyman Memorial Lecture, Robbins (1983) has outlined a wide class of problems concerning the general empirical Bayes approach and the linear empirical Bayes approach to estimation. In this paper we shall study a special case which includes several important standard distributions. Specifically let (θ, x) be a random vector such that θ has a distribution function G , and the conditional expectation of x given θ satisfies

$$E(x|\theta) = \theta. \quad (1.1)$$

Suppose it is desired to use a linear function $A+Bx$ of the observed x to estimate the unknown parameter θ . If the loss function is squared error, the best linear estimate is

$$t(x) = E\theta + \frac{\text{Cov}(\theta, x)}{\text{Var } x} (x - Ex). \quad (1.2)$$

and the mean squared error is

$$E(t-\theta)^2 = \text{Var } \theta - \frac{\text{Cov}^2(\theta, x)}{\text{Var } x}. \quad (1.3)$$

Assume, in addition to (1.1), that

$$\text{Var}(x|\theta) = a + b\theta + c\theta^2. \quad (1.4)$$

for some known constants a, b , and c . Then (1.2) can be written as

$$t(x) = Ex + (1 - \frac{c \text{Var } x + a + b Ex + c E^2 x}{(c+1) \text{Var } x}) (x - Ex) \quad (1.5)$$

which is computable if Ex and $\text{Var } x$ are known.

We shall be dealing with the case when Ex and $\text{Var } x$ are unknown. However we are faced with a large number n of independent versions of the component problem: $(\theta_1, x_1), \dots, (\theta_n, x_n)$ are independent random vectors having the same distributions as (θ, x) . Robbins (1983) has proposed to estimate Ex and $\text{Var } x$ respectively by

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i, \quad \text{and} \quad s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2, \quad (1.6)$$

and use the following statistic (with $h = cs^2 + a + b\bar{x} + c\bar{x}^2$),

$$\tilde{\theta}_i = \bar{x} + (1 - \frac{h}{(c+1)s^2})^+ (x_i - \bar{x}) \quad (1.7)$$

to estimate θ_i , for each $i=1, \dots, n$. He has also hoped that under some mild restrictions on the nature of G , with some reasonable rapidity as n tends to infinity

$$E(\tilde{\theta}_i - \theta_i)^2 \rightarrow E(t - \theta)^2. \quad (1.8)$$

We shall assume that the best linear estimate (1.2) is also the best general estimate $E(\theta|x)$. This assumption will reduce the class of possible distributions for θ . For instance, if x has a distribution from an exponential family with parameter θ , then the above assumption will limit the class of the prior distributions to the conjugate family. See Diaconis and Ylvisaker (1979). However, even this special case will be wide enough to include many standard distributions used in practice. In this case we shall verify (1.8). Indeed we shall consider the cumulative regret

$$R_n = \sum_{i=1}^n (E(\tilde{\theta}_i - \theta_i)^2 - E(t(x_i) - \theta_i)^2) \quad (1.9)$$

of using $\tilde{\theta}_i$ instead of $t(x_i)$ because of the statistician's ignorance of $E x$ and $\text{Var } x$. It can be shown that even as n goes to infinity R_n remains bounded so that (1.8) will hold. We summarize the main results in the following theorem and leave the proof to the next section.

Theorem 1. Let (θ, x) , (θ_1, x_1) , ... be independent and identically distributed nondegenerate random vectors such that

- (i) $E(x|\theta) = \theta$, $\text{Var}(x|\theta) = a + b\theta + c\theta^2$.
- (ii) $E(\theta|x) = E\theta + (x - Ex) \text{Cov}(\theta, x) / \text{Var } x$.
- (iii) $E x^6 < \infty$.

For each $n=2, \dots$, and for each $i=1, 2, \dots, n$, define

$$\begin{aligned}\bar{x} &= \bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{and} \quad s^2 = s_n^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2, \\ t_i &= Ex + (1 - \frac{c \text{Var } x + a + b Ex + c E^2 x}{(c+1) \text{Var } x}) (x_i - Ex), \\ \tilde{\theta}_i &= \bar{x} + (1 - \frac{cs^2 + a + b\bar{x} + c\bar{x}^2}{(c+1)s^2})^+ (x_i - \bar{x}), \quad \text{and} \\ R_n &= \sum_{i=1}^n (E(\tilde{\theta}_i - \theta_i)^2 - E(t_i - \theta_i)^2).\end{aligned}\tag{1.10}$$

Then

$$\lim_{n \rightarrow \infty} R_n = \frac{H^2}{(c+1)^2 \gamma^2} + \frac{\sigma^2}{(c+1)^2 \gamma^6}\tag{1.11}$$

where $\mu = Ex$, $\gamma^2 = \text{Var } x$, $\mu_3 = E(x-\mu)^3$, $\mu_4 = E(x-\mu)^4$,

$$\begin{aligned}H &= c\gamma^2 + a + b\mu + c\mu^2, \quad \text{and} \\ \sigma^2 &= (\mu_4 - \gamma^4)(a + b\mu + c\mu^2)^2 + \gamma^6(b + 2c\mu)^2 \\ &\quad - 2\gamma^2\mu_3(a + b\mu + c\mu^2)(b + 2c\mu)\end{aligned}\tag{1.12}$$

For the special case when $b=c=0$, a slightly more general result can be established under weaker conditions. The result is in Theorem 2.

Theorem 2. Let $(\theta_1, x_1), (\theta_2, x_2), \dots$ be independent random vectors satisfying the following conditions:

(A) For all i ,

- (i) $Ex_i = \mu$ and $\text{Var } x_i = \gamma^2 > 0$.
- (ii) $E(x_i | \theta_i) = \theta_i$ and $\text{Var}(x_i | \theta_i) = a < \gamma^2$.
- (iii) $E(\theta_i | x_i) = E\theta_i + (x_i - \mu) \text{Cov}(\theta_i, x_i) / \gamma^2$.

(B) (iv) $\{(x_i - \mu)^2, i \geq 1\}$ satisfies the Lindeberg condition.

$$(v) \quad \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2 \text{ converges in probability to } \gamma^2, \\ \text{and } \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \text{Var}(x_i - \mu)^2 \text{ is finite.}$$

For each $n = 2, \dots$, and each $i = 1, 2, \dots, n$, put

$$\bar{x} = \bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i, \quad s^2 = s_n^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2, \quad (1.13)$$

$$t_i = Ex_i + (1 - \frac{a}{\text{Var } x_i})(x_i - Ex_i),$$

$$\tilde{\theta}_i = \bar{x} + (1 - \frac{a}{s^2})(x_i - \bar{x}), \quad \text{and}$$

$$R_n = \sum_{i=1}^n (E(\tilde{\theta}_i - \theta_i)^2 - E(t_i - \theta_i)^2).$$

Then

$$\lim_{n \rightarrow \infty} R_n = \frac{a^2}{\gamma^2} + \frac{a^2}{\gamma^6} K, \quad (1.14)$$

where $K = \lim_{n \rightarrow \infty} \frac{1}{n} V_n$, and $V_n = \sum_{i=1}^n \text{Var}(x_i - \mu)^2$.

2. PROOF OF THE MAIN RESULTS

We need some preliminary results for Theorem 1.

Lemma 1. Let x, x_1, x_2, \dots , be independent and identically distributed random variables with $Ex^4 < \infty$. Let the following notation be used: $\mu = Ex$, $\gamma^2 = \text{Var } x$, $\mu_3 = E(x - \mu)^3$ and $\mu_4 = E(x - \mu)^4$.

For each $n \geq 1$, put

$$W_{1n} = \frac{1}{\sqrt{n}} \left(\sum_{i=1}^n (x_i - \mu)^2 - \gamma^2 \right), \quad (2.1)$$

$$W_{2n} = \frac{1}{\sqrt{n}} \left(\sum_{i=1}^n \mu(x_i - \mu)^2 - \gamma^2 x_i \right), \quad \text{and}$$

$$W_{3n} = \sqrt{n} (\bar{x}^2 - \mu^2).$$

Then as n tends to infinity, (W_{1n}, W_{2n}, W_{3n}) converges in distribution to a multivariate normal distribution with mean $\underline{0}$ and

covariance matrix $\Sigma = (\sigma_{ij})$, where

$$\begin{aligned}\sigma_{11} &= \mu_4 - \gamma^4, \quad \sigma_{22} = \mu^2(\mu_4 - \gamma^4) + \gamma^6 - 2\gamma^2\mu\mu_3, \\ \sigma_{33} &= 4\mu^2\gamma^2, \quad \sigma_{12} = \mu(\mu_4 - \gamma^4) - \gamma^2\mu_3, \\ \sigma_{13} &= 2\mu\mu_3, \quad \text{and} \quad \sigma_{23} = 2\mu^2\mu_3 - 2\mu\gamma^4.\end{aligned}\quad (2.2)$$

The proof of the Lemma is straightforward and is omitted.

Corollary 1. Under the same conditions as Lemma 1, as $n \rightarrow \infty$, $(a+c\mu^2)W_{1n} + bW_{2n} - c\gamma^2W_{3n}$ has an asymptotic normal distribution with mean 0 and variance

$$\begin{aligned}\sigma^2 &= (\mu_4 - \gamma^4)(a+b\mu+c\mu^2)^2 + \gamma^6(b+2c\mu)^2 \\ &\quad - 2\gamma^2\mu_3(a+b\mu+c\mu^2)(b+2c\mu).\end{aligned}\quad (2.3)$$

Lemma 2. Let x, x_1, \dots , be independent and identically distributed random variables with mean μ and variance

$$\gamma^2 = a + bEx + cE^2x + d, \quad (2.4)$$

where a, b, c and d are constants and $d > 0$. Assume $Ex^4 < \infty$.

For each $n \geq 2$, put

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{and} \quad s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2. \quad (2.5)$$

Then as n tends to infinity

$$nP[s^2 \leq a+b\bar{x}+c\bar{x}^2] \rightarrow 0. \quad (2.6)$$

Proof: Choose $\delta > 0$, such that $\epsilon = d - |c+1|\delta - |b+2\mu c|\sqrt{\delta} > 0$.

Then

$$\begin{aligned}P[s^2 \leq a+b\bar{x}+c\bar{x}^2] & \\ \leq P[\Sigma(x_i - \mu)^2 - \gamma^2 \leq na + nb\bar{x} + nc\bar{x}^2 - n\gamma^2 + n(\bar{x} - \mu)^2] & \\ \leq P[\Sigma(x_i - \mu)^2 - \gamma^2 \leq -\epsilon n, (\bar{x} - \mu)^2 < \delta] & \\ + P[(\bar{x} - \mu)^2 \geq \delta]. &\end{aligned}\quad (2.7)$$

Let $B = \{\sum (x_i - \mu)^2 - \gamma^2 \leq -\epsilon n\}$. Then the first term in (2.7) is less than or equal to

$$P[|\sum (x_i - \mu)^2 - \gamma^2| \geq \epsilon n] \quad (2.8)$$

$$\leq \frac{1}{\epsilon^2 n^2} \int_B (\sum (x_i - \mu)^2 - \gamma^2)^2 dP$$

which is $o(1/n)$ as n tends to infinity by the uniform integrability of $\{(\sum (x_i - \mu)^2 - \gamma^2)^2/n, n \geq 1\}$ implied by $Ex^4 < \infty$. For the second term in (2.7)

$$P[(\bar{x} - \mu)^2 \geq \delta] \quad (2.9)$$

$$\leq \frac{1}{\delta^2 n^4} (n(E(x - \mu)^4 - \gamma^4) + (3n^2 - 2n)\gamma^4)$$

which is $o(1/n)$ as n tends to infinity. This concludes the proof of Lemma 2.

Lemma 3. Let x, x_1, \dots be independent and identically distributed random variables with mean μ and variance γ^2 . Assume that $Ex^6 < \infty$. For each $n \geq 2$, put

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{and} \quad s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2. \quad (2.10)$$

Then the following families of random variables are uniformly integrable:

$$\begin{aligned} (i) & \quad \{ns^2(s^2 - \gamma^2)^2, n \geq 2\}, \\ (ii) & \quad \{ns^2(\bar{x} - \mu)^2, n \geq 2\}, \\ (iii) & \quad \{ns^2(\bar{x}^2 - \mu^2)^2, n \geq 2\}. \end{aligned} \quad (2.11)$$

Proof. We shall verify (i) and (iii). The verification for (ii) is entirely analogous and hence omitted. For (i),

$$\begin{aligned} & ns^2(s^2 - \gamma^2)^2 \quad (2.12) \\ &= n(s^2 - \gamma^2)^3 + n\gamma^2(s^2 - \gamma^2)^2 \\ &\leq \frac{K}{n^2} |\sum (x_i - \mu)^2 - \gamma^2|^3 + \frac{K}{n^2} |\gamma^2 - n(\bar{x} - \mu)^2|^3 \\ &\quad + \frac{K\gamma^2}{n} |\sum (x_i - \mu)^2 - \gamma^2|^2 + \frac{K\gamma^2}{n} |\gamma^2 - n(\bar{x} - \mu)^2|^2, \end{aligned}$$

for some constant K .

Since $E x^6 < \infty$, it is clear the four terms on the bottom line of (2.12) are uniformly integrable. For (iii), for any event D with $P[D]$ small,

$$E n s^2 (\bar{x}^2 - \mu^2)^2 I_D \quad (2.13)$$

$$\leq K \left(E \left(\frac{\sum (x_i - \mu)^2}{n} \right)^3 I_D \right)^{1/3} \left(E (n(\bar{x}^2 - \mu^2)^2)^{3/2} I_D \right)^{2/3},$$

which can be made small uniformly in n .

Now we are ready to give the proofs. That the convergence in distribution of random variables together with uniform integrability implies moment convergence is used. For a reference, see (Chow and Teicher (1978), Section 8.1).

Proof of Theorem 1. Let

$$h = c s^2 + a + b \bar{x} + c \bar{x}^2. \quad (2.14)$$

From the identity

$$\begin{aligned} (\tilde{\theta}_i - \theta_i)^2 - (t_i - \theta_i)^2 \\ = (\tilde{\theta}_i - t_i)^2 + 2(\tilde{\theta}_i - t_i)(t_i - \theta_i); \end{aligned} \quad (2.15)$$

taking expectation and summation and by assumptions (i) and (ii) we have the cumulative regret

$$R_n = \frac{H^2}{(c+1)^2 \gamma^2} + E \left(\frac{H}{(c+1)\gamma^2} - \frac{h}{\max(h, (c+1)s^2)} \right)^2 \sum (x_i - \bar{x})^2 \quad (2.16)$$

As n tends to infinity s^2 and h will go to γ^2 and H respectively with probability one. Since, (see Robbins (1983)),

$$\gamma^2 = \frac{H}{c+1} + \text{Var } \theta, \quad (2.17)$$

and θ is nondegenerate, asymptotically the term inside the expectation sign in (2.16) is equivalent to

$$\frac{n}{(c+1)^2 \gamma^4} \frac{(s^2 H - \gamma^2 h)^2}{s^2} \quad (2.18)$$

$$\begin{aligned} &= \frac{1}{(c+1)^2 \gamma^4 s^2} ((a+c\mu^2) \sqrt{n}(s^2 - \gamma^2) + b\sqrt{n}(\mu s^2 - \gamma^2 \bar{x}) - c\gamma^2 \sqrt{n}(\bar{x}^2 - \mu^2))^2 \\ &= \frac{1}{(c+1)^2 \gamma^6} ((a+c\mu^2) \frac{\sum (x_i - \mu)^2 - \gamma^2}{\sqrt{n}} + b \frac{\sum \mu(x_i - \mu)^2 - \gamma^2 x_i}{\sqrt{n}} \\ &\quad - c\gamma^2 \sqrt{n}(\bar{x}^2 - \mu^2) + o_p(1))^2. \end{aligned}$$

By Corollary 1 and (2.18), $(\chi_1^2$ denoting chi-squared with 1 d.f.),

$$T = \left(\frac{H}{(c+1)\gamma^2} - \frac{h}{\max(h, (c+1)s^2)} \right)^2 \sum (x_i - \bar{x})^2 \quad (2.19)$$

converges in distribution to $\frac{\sigma^2}{(c+1)^2 \gamma^6} \chi_1^2$, where σ^2 is defined

in (2.3) Next we shall show that $\{T, n \geq 2\}$ is uniformly integrable so that as n tends to infinity

$$ET \rightarrow \frac{\sigma^2}{(c+1)^2 \gamma^6}. \quad (2.20)$$

Let A be the event $\{(c+1)s^2 \leq h\}$, then for some positive K

$$\begin{aligned} ET \mathbb{I}_A &\leq K \int_A (na + nb\bar{x} + nc\bar{x}^2) dP \\ &\leq K(a+b\mu+c\mu^2)nP[A] + Kbn \int_A |\bar{x} - \mu| dP + Kcn \int_A |\bar{x}^2 - \mu^2| dP \\ &\leq K(a+b\mu+c\mu^2)nP[A] + Kb\sqrt{n} ((En(\bar{x} - \mu)^2) P[A])^{\frac{1}{2}} \\ &\quad + Kc\sqrt{n} (En(\bar{x}^2 - \mu^2)^2 P[A])^{\frac{1}{2}}, \end{aligned} \quad (2.21)$$

which is $o(1)$ by Lemma 2. On the complement of A ,

$$\begin{aligned} &\frac{(c+1)^2 \gamma^4}{2} T \\ &= \frac{(c+1)^2 \gamma^4}{2} \left(\frac{H}{(c+1)\gamma^2} - \frac{h}{(c+1)\gamma^2} + \frac{h}{(c+1)\gamma^2} - \frac{h}{(c+1)s^2} \right)^2 \sum (x_i - \bar{x})^2 \end{aligned} \quad (2.22)$$

$$\begin{aligned}
&\leq (H-h)^2 \Sigma(x_i - \bar{x})^2 + \frac{nh^2}{s^2} (s^2 - \gamma^2)^2 \\
&\leq 4c^2 (s^2 - \gamma^2)^2 \Sigma(x_i - \bar{x})^2 + 4b^2 (\bar{x} - \mu)^2 \Sigma(x_i - \bar{x})^2 \\
&\quad + 4c^2 (\bar{x}^2 - \mu^2)^2 \Sigma(x_i - \bar{x})^2 + (c+1)^2 ns^2 (s^2 - \gamma^2)^2.
\end{aligned}$$

By Lemma 3, the four terms on the bottom line of (2.22) are uniformly integrable. Therefore we have the regret

$$\lim_{n \rightarrow \infty} R_n = \frac{H^2}{(c+1)^2 \gamma^2} + \frac{\sigma^2}{(c+1)^2 \gamma^6}. \quad (2.23)$$

Remark. The expression for σ^2 contains terms up to the fourth moment. Although it has terms of the eighth order (e.g. $c^2 \mu^4 \mu_4$), the sixth moment assumption is to ensure the uniform integrability of $\{(\frac{1}{\sqrt{n}} \Sigma((x_i - \mu)^2 - \gamma^2))^3, n \geq 1\}$ in Lemma 3. Nevertheless it is reasonable to conjecture that condition (iii) in Theorem 1 can be replaced by $Ex^4 < \infty$.

Proof of Theorem 2. From the identity

$$\begin{aligned}
(\tilde{\theta}_i - \theta_i)^2 - (t_i - \theta_i)^2 &= (\tilde{\theta}_i - t_i)^2 \\
&\quad + 2(\tilde{\theta}_i - t_i)(t_i - \theta_i);
\end{aligned} \quad (2.24)$$

taking summation and expectation and by assumptions (i), (ii), and (iii), and definition (1.13) we have the regret equal to

$$R_n = \frac{a^2}{\gamma^2} + a^2 E\left(\frac{1}{\gamma^2} - \frac{1}{\max(s^2, a)}\right)^2 \Sigma(x_i - \bar{x})^2. \quad (2.25)$$

Let $v_i = \text{Var}(x_i - \mu)^2$ and $V_n = v_1 + v_2 + \dots + v_n$. Consider the event $A = \{s^2 \leq a\}$, then

$$\begin{aligned}
&E\left(\frac{1}{\gamma^2} - \frac{1}{\max(x^2, a)}\right)^2 \Sigma(x_i - \bar{x})^2 I_A \\
&\leq cn P\{s^2 \leq a\}
\end{aligned} \quad (2.26)$$

for some positive constant c . And

$$\begin{aligned}
P[s^2 \leq a] &\leq P[\Sigma(x_i - \mu)^2 - \gamma^2 \leq (a - \gamma^2)n + n(\bar{x} - \mu)^2] \\
&\leq P[\Sigma(x_i - \mu)^2 - \gamma^2 \leq (a - \gamma^2)n + n(\bar{x} - \mu)^2, (\bar{x} - \mu)^2 < \delta] \\
&\quad + P[(\bar{x} - \mu)^2 \geq \delta].
\end{aligned} \tag{2.27}$$

Choose $\delta > 0$ such that $a - \gamma^2 + \delta = -\varepsilon < 0$. Let B be

$$\{\Sigma(x_i - \mu)^2 - \gamma^2 \leq -\varepsilon n\},$$

then

$$\begin{aligned}
&nP[\Sigma(x_i - \mu)^2 - \gamma^2 \leq -\varepsilon n] \\
&\leq \frac{V_n}{\varepsilon^2 n} \int_B \frac{(\Sigma(x_i - \mu)^2 - \gamma^2)^2}{V_n} dP
\end{aligned} \tag{2.28}$$

which goes to zero as n tends to infinity by conditions (iv) and (v), and Brown's Theorem (see Chow and Teicher (1978), p. 398).

Next, consider

$$\begin{aligned}
&nP[(\bar{x} - \mu)^2 \geq \delta] \\
&\leq \frac{n}{\delta^2} E(\bar{x} - \mu)^4 = \frac{1}{\delta^2 n^3} (\Sigma \text{Var}(x_i - \mu)^2 + (3n^2 - 2n)\gamma^4)
\end{aligned} \tag{2.29}$$

which goes to zero as n tends to infinity by condition (v). On the complement of A , and for any event D

$$\begin{aligned}
&E\left(\frac{1}{\gamma^2} - \frac{1}{\max(s^2, a)}\right)^2 \Sigma(x_i - \bar{x})^2 I_A^c I_D \\
&= E\left(\frac{\Sigma(x_i - \bar{x})^2 - (n-1)\gamma^2}{\gamma^2 \Sigma(x_i - \bar{x})^2}\right)^2 \Sigma(x_i - \bar{x})^2 I_A^c I_D \\
&\leq \frac{V_n}{(n-1)a\gamma^4} E \frac{(\Sigma(x_i - \bar{x})^2 - (n-1)\gamma^2)^2}{V_n} I_A^c I_D
\end{aligned} \tag{2.30}$$

which may be made very small uniformly in n if $P[D]$ is small for the same reason as in (2.28). Therefore the family

$$\left\{ \left(\frac{1}{\gamma^2} - \frac{1}{\max(s^2, a)} \right)^2 \sum (x_i - \bar{x})^2, n \geq 2 \right\} \quad (2.31)$$

is uniformly integrable. And by condition (v), as n tends to infinity s^2 tends to γ^2 and \bar{x} tends to μ in probability. By condition (iv), as n tends to infinity

$$\frac{\sum (x_i - \bar{x})^2 - (n-1)\gamma^2}{\sqrt{V_n}} \rightarrow N(0,1) \text{ in distribution.} \quad (2.32)$$

Hence

$$E \left(\frac{1}{\gamma^2} - \frac{1}{\max(s^2, a)} \right)^2 \sum (x_i - \bar{x})^2 \rightarrow \frac{1}{\gamma^6} \lim_{n \rightarrow \infty} \frac{V_n}{n}; \quad (2.33)$$

that is

$$\lim_{n \rightarrow \infty} R_n = \frac{a^2}{\gamma^2} + \frac{a^2}{\gamma^6} \lim_{n \rightarrow \infty} \frac{V_n}{n}. \quad (2.34)$$

Corollary: Let $(\theta_1, x_1), (\theta_2, x_2), \dots$ be independent and identically distributed random vectors satisfying condition (A) in the Theorem. If $E x^4 < \infty$, R_n and $\tilde{\theta}_i$ are defined in the same way as in (1.13) and (1.14), then

$$\lim_{n \rightarrow \infty} R_n = \frac{a^2}{\gamma^2} E \left(\frac{x - \mu}{\gamma} \right)^4. \quad (2.35)$$

Example 1. Suppose θ has a common normal distribution with mean μ and variance τ^2 and given θ , x has a normal distribution with mean θ and variance a . Then x has a normal distribution with mean μ and variance $\gamma^2 = a + \tau^2$. Obviously the conditions of the corollary hold; hence the regret R_n is $3a^2/(a + \tau^2) + o(1)$, as n tends to infinity.

$\tilde{\theta}_i$ in this normal case is a variant of the James-Stein estimator, which has been studied extensively in the literature. See Efron and Morris (1973).

Example 2. Suppose θ has an inverted gamma distribution with density function

$$g(\theta) = \begin{cases} \left(\frac{\beta}{\theta}\right)^{\alpha+1} \frac{e^{-\beta/\theta}}{\Gamma(\alpha)} \frac{1}{\beta} & \text{if } \theta > 0 \\ 0 & \text{if } \theta \leq 0, \end{cases}$$

where $\beta > 0$ and $\alpha > 6$, and given θ , x has an exponential distribution with mean θ and variance θ^2 . The conditions (i), (ii) and (iii) in Theorem 1 hold with $a=b=0$ and $c=1$. It can be computed that, for any $0 \leq p \leq 6$;

$$Ex^p = \frac{\beta^p \Gamma(p+1) \Gamma(\alpha-p)}{\Gamma(\alpha)}$$

If the linear empirical Bayes estimators are used, then the cumulative regret R_n will satisfy

$$\lim_{n \rightarrow \infty} R_n = \frac{2(\alpha-1)(\alpha^2-4\alpha+6)\beta^2}{\alpha^2(\alpha-2)(\alpha-3)(\alpha-4)}.$$

Example 3. Suppose θ has a gamma distribution with the density function

$$g(\theta) = \begin{cases} 0 & \text{if } \theta \leq 0 \\ \frac{\beta^\alpha \theta^{\alpha-1} e^{-\beta\theta}}{\Gamma(\alpha)} & \text{if } \theta > 0, \end{cases}$$

where α and β are positive constants, and given θ , x has a Poisson distribution with mean θ . In this case, the conditions (i), (ii) and (iii) in Theorem 1 hold with $a=c=0$ and $b=1$. If the linear empirical Bayes estimators are used, then the regret R_n will satisfy

$$\lim_{n \rightarrow \infty} R_n = \frac{3\alpha(\beta+1) + 2\beta+3}{(\beta+1)^2}.$$

ACKNOWLEDGEMENT

The author thanks the referees for invaluable suggestions. The research is supported by a University of South Carolina Research and Productive Scholarship and the United States Air Force Office of Scientific Research and Army Research Office under Grant No. AFOSR-84-0156.

BIBLIOGRAPHY

- Chow, Y. S. and Teicher, H. (1978). Probability Theory, Springer Verlag.
- Diaconis, P. and Ylvisaker, D. (1979). Conjugate priors for exponential families. Annals of Statistics, 7, 269-281.
- Efron, B. and Morris, C. (1973). Stein's estimation rule and its competitors - an empirical Bayes approach. Journal of American Statistical Association, 68, 117-130.
- Robbins, H. (1983). Some thoughts on empirical Bayes estimation. Jerzy Newman Memorial Lecture. Annals of Statistics, 11, 713-723.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED		REPORT DOCUMENTATION PAGE	
1a. REPORT SECURITY CLASSIFICATION		1d. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release, distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Statistics Technical Report No. 112		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Department of Statistics	8b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research	
6c. ADDRESS (City, State and ZIP Code) University of South Carolina Columbia, SC 29208		7b. ADDRESS (City, State and ZIP Code) Directorate of Mathematical & Information Sciences, Bolling AFB, DC 20332	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR, ARO	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-84-0156	
8c. ADDRESS (City, State and ZIP Code) Bolling AFB, DC 20332		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304
		TASK NO. A5	WORK UNIT NO.
11. TITLE (Include Security Classification) On the bounded regret of empirical Bayes estimators.			
12. PERSONAL AUTHOR(S) K. F. Yu			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Yr., Mo., Day) May, 1985	15. PAGE COUNT 14
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
		Empirical Bayes; bounded regret; uniform integrability.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Let $(\theta_1, x_1), \dots, (\theta_n, x_n)$ be independent and identically distributed random vectors with $E(x \theta) = \theta$ and $\text{Var}(x \theta) = a + b\theta + c\theta^2$. Let t_i be the linear Bayes estimator of θ_i and $\tilde{\theta}_i$ be the linear empirical Bayes estimator of θ_i as proposed in Robbins (1983), when $E x$ and $\text{Var } x$ are unknown to the statistician. The regret of using $\tilde{\theta}_i$ instead of t_i because of ignorance of the mean and the variance is $r_i = E(\tilde{\theta}_i - \theta_i)^2 - E(t_i - \theta_i)^2$. Under appropriate conditions cumulative regret $R_n = r_1 + \dots + r_n$ is shown to have a finite limit even when n tends to infinity. The limit can be explicitly computed in terms of a, b, c and the first four moments of x .			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Maj. Brian W. Woodruff		22b. TELEPHONE NUMBER (Include Area Code) (202) 765-5027	22c. OFFICE SYMBOL NM

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Statistics Technical Report No. 112	2. GOVT ACCESSION NO. N/A	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle) On the bounded regret of empirical Bayes estimators		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER Stat.Tech.Rpt. No. 112
7. AUTHOR(s) K. F. Yu		8. CONTRACT OR GRANT NUMBER(s) MIPR ARO 139-85
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics University of South Carolina Columbia, SC 29208		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102 F 2304A5 N/A
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE May 1985
		13. NUMBER OF PAGES 14
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Empirical Bayes; bounded regret; uniform integrability.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let $(\theta_1, x_1), \dots, (\theta_n, x_n)$ be independent and identically distributed random vectors with $E(x \theta) = \theta$ and $\text{Var}(x \theta) = a + b\theta + c\theta^2$. Let t_i be the linear Bayes estimator of θ_i and $\tilde{\theta}_i$ be the linear empirical Bayes Estimator of θ_i as proposed in Robbins (1983), when EX and $\text{Var } x$ are unknown to the statistician. The regret of using $\tilde{\theta}_i$ instead of t_i because of ignorance of the mean and the		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

variance is $r_i = E(\tilde{\theta}_i - \theta_i)^2 - E(t_i - \theta_i)^2$. Under appropriate conditions cumulative regret $R_n = r_1 + \dots + r_n$ is shown to have a finite limit even when n tends to infinity. The limit can be explicitly computed in terms of a, b, c and the first four moments of x .

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

END

DTIC

7-86